

Optimality Conditions for Fractional Minmax Programming

CHANCHAL SINGH

Department of Mathematics, St. Lawrence University, Canton, New York 13617

Submitted by L. Zadeh

1. INTRODUCTION

The purpose of this paper is to generalize the optimality conditions of Schmitendorf [6] and of fractional programming studied by Aggarwal and Saxena [1]. Aggarwal and Saxena considered the following problem:

$$\begin{aligned} &\text{Minimize } F(x) = N(x)/D(x) = (f(x) + (x^t B x)^{1/2})/h(x) \\ &\text{subject to } g(x) \leq 0. \end{aligned}$$

Where $x \in R^n$, B is an $n \times n$ symmetric positive semidefinite matrix, $f(\cdot)$, $-h(\cdot)$ are real valued, convex, differentiable functions defined on R^n and $g(\cdot)$ is a p -dimensional, convex, differentiable function also defined on R^n . The function $h(x) > 0$ for all feasible x .

We consider the following generalization of the Aggarwal-Saxena problem:

$$\begin{aligned} &\text{Minimize } F(x) = \sup_{y \in Y} (f(x, y) + (x^t B x)^{1/2})/h(x, y) \\ &\text{subject to } g(x) \leq 0 \end{aligned} \tag{1}$$

where Y is a compact subset of R^m , $f(\cdot, \cdot): R^n \times R^m \rightarrow R$ is C^1 on $R^n \times R^m$ and $g(\cdot): R^n \rightarrow R^p$ is C^1 on R^n ; B is an $n \times n$ positive semidefinite matrix; $h(\cdot, \cdot): R^n \times R^m \rightarrow R$ is C^1 on $R^n \times R^m$. Throughout the paper we assume that $h(x, y) > 0$ for each (x, y) in $X \times Y$, where X is the set of feasible solutions of Problem (1), i.e., $X = \{x \in R^n: g(x) \leq 0\}$.

The necessary and sufficient conditions to be established are based on the optimality conditions developed by Schmitendorf [6] for a static minmax problem. This application is possible when the functions involved are differentiable at an optimal point or a candidate for optimal point. However, differentiability conditions may not hold due to the presence of $(x^t B x)^{1/2}$ in the objective function. In this situation, the constraint qualification used by Aggarwal and Saxena and introduced by Mond [4] is extended for our case.

As pointed out by Mond and Schechter [5], Mond's constraint qualification leads to the usual Kuhn-Tucker constraint qualification of classical nonlinear programming.

2. PRELIMINARIES

We let X , the set of feasible solutions, be a compact set.

$$\begin{aligned} I(x^*) &= \{i: g_i(x^*) = 0\} \\ Y(x) &= \{y \in Y; (f(x, y) + (x^t Bx)^{1/2})/h(x, y) \\ &= \sup_{z \in Y} (f(x, z) + (x^t Bx)^{1/2})/h(x, z)\} \end{aligned}$$

K = the set of triplets (s, t, \bar{y}) , where s ranges over the positive integers such that $1 \leq s \leq n+1$; $t = (t_1, \dots, t_s)$ an s -dimensional vector with $t_i \geq 0$, $\sum_{i=1}^s t_i = 1$; $\bar{y} = (y_1, \dots, y_s)$ an ms -dimensional vector with $y_i \in Y(x)$ ($i = 1, \dots, s$) for some $x \in R^n$.

For $x_0 \in X$, $(s, t, \bar{y}_0) \in K$, $y_0^i \in Y(x_0)$,

$$v_0 = (f(x_0, y_0^i) + (x_0^t Bx_0)^{1/2})/h(x_0, y_0^i),$$

we let

$$Q_{\bar{y}}(x_0) = \{z: z^t \nabla g_i(x_0) \leq 0 \quad i \in I(x_0)\}$$

and

$$\begin{aligned} & z^t \left(\sum_{i=1}^s (t_i/h(x_0, y_0^i)) (\nabla_x f(x_0, y_0^i) - \nabla_x v_0 \right. \\ & \quad \left. \cdot h(x_0, y_0^i)) \right) + z^t Bx_0 / (x_0^t Bx_0)^{1/2} < 0 \quad \text{if } x_0^t Bx_0 > 0; \\ & z^t \left(\sum_{i=1}^s (t_i/h(x_0, y_0^i)) (\nabla_x f(x_0, y_0^i) - \nabla_x v_0 \right. \\ & \quad \left. \cdot h(x_0, y_0^i)) \right) + (z^t Bz)^{1/2} < 0 \quad \text{if } x_0^t Bx_0 = 0 \Big\}. \end{aligned}$$

We state the following two lemmas of Eisenberg [3] for easy reference.

LEMMA 1 [3]. *Let B be a symmetric positive semidefinite matrix. Then $x^t Bz \leq (x^t Bx)^{1/2} (z^t Bz)^{1/2}$ for each x and z .*

LEMMA 2 [3]. Let A be an $m \times n$ matrix and H be a symmetric positive semidefinite matrix. Let p be an n -dimensional vector. Then there exists an x such that $Ax \geq 0 \Rightarrow p^t x + (x^t H x)^{1/2} \geq 0$ if and only if there exist $u \in R^m$, $u \geq 0$, $w \in R^n$ such that $A^t u = Hw + p$, $Aw \geq 0$, $w^t H w \leq 1$.

The next lemma is an extension of a result of Bhatia.

LEMMA [2]. Suppose $x_0 \in S$, a compact subset of R^n . Let $y_0 \in Y(x_0)$. Then x_0 minimizes $(f(x, y_0) + (x^t B x)^{1/2})/h(x, y_0)$ if and only if it minimizes $(f(x, y_0) + (x^t B x)^{1/2}) - v_0 h(x, y_0)$, where $v_0 = \text{Min}_{x \in S} (f(x, y_0) + (x^t B x)^{1/2})/h(x, y_0)$.

Proof. Suppose x_0 minimizes $(f(x, y_0) + (x^t B x)^{1/2})/h(x, y_0)$ but it does not minimize $f(x, y_0) + (x^t B x)^{1/2} - v_0 h(x, y_0)$. Let x_1 be such that it minimizes $f(x, y_0) + (x^t B x)^{1/2} - v_0 h(x, y_0)$. Then $f(x_1, y_0) + (x_1^t B x_1)^{1/2} - v_0 h(x_1, y_0) < f(x_0, y_0) + (x_0^t B x_0)^{1/2} - v_0 h(x_0, y_0) = 0$. Therefore $(f(x_1, y_0) + (x_1^t B x_1)^{1/2})/h(x_1, y_0) < v_0$. This means that v_0 is not the minimum value, which contradicts the hypothesis that it is. Hence x_0 also minimizes $f(x, y_0) + (x^t B x)^{1/2} - v_0 h(x, y_0)$.

Conversely suppose x_0 minimizes $f(x, y_0) + (x^t B x)^{1/2} - v_0 h(x, y_0)$ over S . But x_0 certainly minimizes $f(x, y_0) + (x^t B x)^{1/2}/h(x, y_0)$ with its value equal to v_0 . So the proof is complete.

3. OPTIMALITY CONDITIONS

The two theorems we are about to establish are direct generalizations of necessary and sufficient conditions of Schmitendorf [6]. (When $B = 0$, $h(x, y) = 1$, our theorems are the same as Schmitendorf's.) These results also generalize Theorems 4.1 and 4.2 of Aggarwal and Saxena [1] because of the presence of vector y in the objective function.

THEOREM 1 (Necessary Conditions). If x_0 is an optimal solution of Problem 1 and $Q_{\bar{y}}(x_0)$ is empty, then there exist a positive integer s , $1 \leq s \leq n + 1$, real vectors $t = (t_1, \dots, t_s)$, $t \geq 0$, $u = (u_1, \dots, u_p)$, $u \geq 0$, $w = (w_1, \dots, w_n)$, $y_0^i \in Y(x_0)$ ($i = 1, \dots, s$) such that

$$\sum_{i=1}^s (t_i/h(x_0, y_0^i))(\nabla_x f(x_0, y_0^i)) + Bw - v_0 \nabla_x h(x_0, y_0) + \sum_{i=1}^p u_i \nabla g_i(x_0) = 0$$

$$u_i g_i(x_0) = 0, i = 1, \dots, p, w^t B w \leq 1, (x_0^t B x_0)^{1/2} = x_0^t B w$$

and

$$\sum_{i=1}^s t_i + \sum_{i=1}^p u_i > 0 \quad \text{where} \quad v_0 = (f(x_0, y_0^i) + (x_0^t B x_0)^{1/2})/h(x_0, y_0^i),$$

$$y_0^i \in Y(x_0) \quad \text{for} \quad i = 1, \dots, s.$$

Proof. Case 1. Suppose $x_0^t B x_0 > 0$. The function $(f(x, y) + (x B x)^{1/2})/h(x, y)$ is differentiable with respect to x for each y . Hence by Schmitendorf's Theorem 1 [6], there exist a positive integer s , vectors $t \in R^s$, $u \in R^p$, $w \in R^n$, $y_0^i \in Y(x_0)$ ($i = 1, \dots, s$) such that $u_i g_i(x_0) = 0$ ($i = 1, \dots, p$), $\sum_{i=1}^s t_i + \sum_{i=1}^p u_i > 0$ and

$$\begin{aligned} & \sum_{i=1}^s ((t_i/h(x_0, y_0^i))(\nabla_x f(x_0, y_0^i) + B x_0)/(x_0^t B x_0)^{1/2}) \\ & - (t_i/(h(x_0, y_0^i))^2)(f(x_0, y_0^i) + (x_0^t B x_0)^{1/2} (\nabla_x h(x_0, y_0^i))) \\ & + \sum_{i=1}^p u_i \nabla g_i(x_0) = 0. \end{aligned} \quad (2)$$

Letting $w = x_0/(x_0^t B x_0)^{1/2}$, (2) reduces to

$$\begin{aligned} & \sum_{i=1}^s (t_i/h(x_0, y_0^i))(\nabla_x f(x_0, y_0^i) + B w - v_0 \nabla_x h(x_0, y_0^i)) \\ & + \sum_{i=1}^p u_i \nabla g_i(x_0) = 0. \end{aligned}$$

Furthermore, since $w = x_0/(x_0^t B x_0)^{1/2}$, it follows that $w^t B w = 1$ and $(x_0^t B x_0)^{1/2} = x_0^t B w$. Therefore in this case all the conclusions of the Theorem hold.

Case 2. $x_0^t B x_0 = 0$. Since $Q_{\bar{y}}(x_0)$ is empty, $z^t \nabla g_i(x_0) \geq 0$ implies

$$z^t \sum_{i=1}^s (t_i/h(x_0, y_0^i))(\nabla_x f(x_0, y_0^i) - \nabla_x v_0 h(x_0, y_0^i)) + (z^t B z)^{1/2} \geq 0.$$

Hence by Lemma 2, there exist $u \geq 0$, $u \in R^{I(x_0)}$, $w \in R^n$ such that

$$u^t \nabla g_{I(x_0)}(x_0) + \sum_{i=1}^s (t_i/h(x_0, y_0^i))(\nabla_x f(x_0, y_0^i) - \nabla_x v_0 h(x_0, y_0^i)) + B w = 0$$

and $w^t B w \leq 1$. Or

$$\sum_{i \in I(x_0)} u_i \nabla g_i(x_0) + \sum_{i=1}^s (t_i/h(x_0, y_0^i))(\nabla_x f(x_0, y_0^i) - \nabla_x v_0 h(x_0, y_0^i)) + B w = 0$$

and $w^t Bw \leq 1$. Since $g_i(x_0) = 0$ for $i \in I(x_0)$, $u_i g_i(x_0) = 0$ for $i \in I(x_0)$. Letting $u_i = 0$ for all i such that $g_i(x_0) < 0$, we have $u_i g_i(x_0) = 0$ ($i = 1, \dots, p$). The fact that $u_i \geq 0$ and $(s, t, \bar{y}_0) \in K$ implies that $\sum_{k=1}^s t_i = 1$ and hence $\sum_{i=1}^s t_i + \sum_{i=1}^p u_i > 0$. From Lemma 1, $x_0^t Bx_0 = 0$ implies $Bx_0 = 0$. Hence $(x_0^t Bx_0)^{1/2} = 0 = x_0^t Bw$. The proof is complete.

THEOREM 2 (Sufficient Conditions). Suppose $g(\cdot)$ is a convex differentiable function, and $f(\cdot, y)$ and $-v_0 h(\cdot, y)$ are convex differentiable functions of x for each $y \in Y$. Suppose there exists a positive integer s , $1 \leq s \leq n+1$, real vectors $t \in R_+^s$, $u \in R_+^p$, $w \in R^n$ and $y_0^i \in Y(x_0)$, for $i = 1, \dots, s$, for some $x_0 \in X$ such that

- (i) $\sum_{i=1}^s (t_i/h(x_0, y_0^i))(\nabla_x f(x_0, y_0^i) - \nabla_x v_0 h(x_0, y_0^i) + Bw) + \sum_{i=1}^p u_i \nabla g_i(x_0) = 0$,
- (ii) $u_i g_i(x_0) = 0$ for $i = 1, \dots, p$,
- (iii) $w^t Bw \leq 1$, $(x_0^t Bx_0)^{1/2} = x_0^t Bw$, $\sum_{i=1}^s t_i > 0$, where $v_0 = (f(x_0, y_0^i) + (x_0^t Bx_0)^{1/2})/h(x_0, y_0^i)$ for $i = 1, \dots, s$. Then x_0 is an optimal solution of Problem (1).

Proof. Suppose x_0 is not an optimal solution of (1). Then there exists an $x_1 \in X$ such that

$$\begin{aligned} & \sup_{y \in Y} ((f(x_1, y) + (x_1^t Bx_1)^{1/2})/h(x_1, y)) \\ & < \sup_{y \in Y} ((f(x_0, y) + (x_0^t Bx_0)^{1/2})/h(x_0, y)). \end{aligned} \quad (3)$$

Also

$$u_i g_i(x_0) = 0, \quad i = 1, \dots, p \quad (4)$$

$$u_i g_i(x_1) \leq 0, \quad i = 1, \dots, p. \quad (5)$$

We note that

$$\begin{aligned} & \sup_{y \in Y} ((f(x_0, y) + (x_0^t Bx_0)^{1/2})/h(x_0, y)) \\ & = (f(x_0, y_0^i) + (x_0^t Bx_0)^{1/2})/h(x_0, y_0^i) \end{aligned} \quad (6)$$

for $i = 1, \dots, s$ and

$$\begin{aligned} & (f(x_1, y_0^i) + (x_1^t Bx_1)^{1/2})/h(x_1, y_0^i) \\ & \leq \sup_{y \in Y} ((f(x_1, y) + (x_1^t Bx_1)^{1/2})/h(x_1, y)). \end{aligned} \quad (7)$$

Hence by (3), (6) and (7),

$$(f(x_1, y_0^i) + (x_1^t B x_1)^{1/2})/h(x_1, y_0^i) < (f(x_0, y_0^i) + (x_0^t B x_0)^{1/2})/h(x_0, y_0^i) \quad (8)$$

for $i = 1, \dots, s$. By hypothesis (i),

$$\begin{aligned} & \sum_{i=1}^s (t_i/h(x_0, y_0^i))(x_1 - x_0)^t (\nabla_x f(x_0, y_0^i) + Bw - v_0 \nabla_x h(x_0, y_0^i)) \\ &= - \sum_{i=1}^p u_i(x_1 - x_0)^t \nabla g_i(x_0). \end{aligned} \quad (9)$$

Since g_i is convex for each i ,

$$\sum_{i=1}^p (u_i g_i(x_1) - u_i g_i(x_0)) \geq \sum_{i=1}^p u_i(x_1 - x_0)^t \nabla g_i(x_0). \quad (10)$$

By (4), (5) and (10), we have

$$\sum_{i=1}^p u_i(x_1 - x_0)^t \nabla g_i(x_0) \leq 0. \quad (11)$$

By (9) and (11),

$$\begin{aligned} & \sum_{i=1}^s (t_i/h(x_0, y_0^i))((x_1 - x_0)^t (\nabla_x f(x_0, y_0^i) - v_0 \nabla_x h(x_0, y_0^i)) \\ &+ x_1^t Bw - x_0^t Bw) \geq 0. \end{aligned} \quad (12)$$

Therefore by the convexity of f and $-h$ with respect to x for each y and using (12),

$$\begin{aligned} & \sum_{i=1}^s (t_i/h(x_0, y_0^i))(f(x_1, y_0^i) - f(x_0, y_0^i) + v_0 h(x_0, y_0^i) \\ &- v_0 h(x_1, y_0^i) + x_1^t Bw - x_0^t Bw) \geq 0. \end{aligned} \quad (13)$$

Since $v_0 = (f(x_0, y_0^i) + (x_0^t B x_0)^{1/2})/h(x_0, y_0^i)$ for $i = 1, \dots, s$ and by Lemma 1, (13) becomes

$$\begin{aligned} & \sum_{i=1}^s (t_i/h(x_0, y_0^i))(f(x_1, y_0^i) + (x_0^t B x_0)^{1/2} - (f(x_0, y_0^i) \\ &+ (x_0^t B x_0)^{1/2})/h(x_0, y_0^i)) \cdot h(x_1, y_0^i) + (x_1^t B x_1)^{1/2} (w^t Bw)^{1/2} - x_0^t Bw \geq 0. \end{aligned} \quad (14)$$

Now by hypothesis (iii), (14) reduces to

$$\sum_{i=1}^s (t_i/h(x_0, y_0^i))(f(x_1, y_0^i) + (x_1^t B x_1)^{1/2} - (f(x_0, y_0^i) + (x_0^t B x_0)^{1/2})/h(x_0, y_0^i)) - h(x_1, y_0^i) \geq 0.$$

Or

$$\sum_{i=1}^s ((t_i/h(x_1, y_0^i)/h(x_0, y_0^i))((f(x_1, y_0^i) + (x_1^t B x_1)^{1/2})/h(x_1, y_0^i)) - (f(x_0, y_0^i) + (x_0^t B x_0)^{1/2})/h(x_0, y_0^i)) \geq 0.$$

But this last inequality is impossible since $t_i \geq 0$, $h(x_0, y_0^i) > 0$, $h(x_1, y_0^i) > 0$ and

$$((f(x_1, y_0^i) + (x_1^t B x_1)^{1/2})/h(x_1, y_0^i)) - (f(x_0, y_0^i) + (x_0^t B x_0)^{1/2})/h(x_0, y_0^i) < 0 \quad \text{by (8).}$$

This contradiction leads to the conclusion that x_0 minimizes Problem (1).

REFERENCES

1. S. P. AGGARWAL AND P. C. SAXENA, A class of fractional functional programming problems, *New Zealand Oper. Res.* **7** (1979), 79–90.
2. D. BHATIA, A note on duality theorem for a nonlinear programming problem, *Management Sci.* **16** (1970), 604–606.
3. E. EISENBERG, Support of a convex function, *Bull. Amer. Math. Soc.* **68** (1962), 192–195.
4. B. MOND, A class of nondifferentiable mathematical programming problems, *J. Math. Anal. Appl.* **46** (1974), 169–174.
5. B. MOND AND M. SCHECHTER, On a constraint qualification in a nondifferentiable programming problem, *Naval Res. Logist. Quart.* **23** (4) (1976), 611–613.
6. W. E. SCHMITENDORF, Necessary conditions and sufficient conditions for static minmax problems, *J. Math. Anal. Appl.* **57** (1977), 683–693.